

Carex globularis

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **3.8**

Moisture indicator value: **7.6**

Disturbance indicator values

Disturbance frequency: **0.14**

Disturbance frequency (herb layer): **1.34**

Disturbance severity: **0.64**

Disturbance severity (herb layer): **0.11**

Mowing frequency: **0.01**

Grazing pressure: **0.18**

Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [BA \(PIC\) Vaccinio-Piceetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T3J Pinus and Larix mire forest](#), [T3K Picea mire forest](#)

Constant species of EUNIS habitats: [T3J Pinus and Larix mire forest](#), [T3K Picea mire](#)

forest

Broad habitat

Occurrence in broad habitats: **Forest**