

Carex mairei

Habitus and growth type

Plant height [m]: **0.4**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.5**

Temperature indicator value: **6.7**

Moisture indicator value: **7.5**

Reaction indicator value: **8**

Nutrient indicator value: **3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.45**

Disturbance frequency (herb layer): **1.91**

Disturbance severity: **0.35**

Disturbance severity (herb layer): **0.33**

Mowing frequency: **1.01**

Grazing pressure: **0.31**

Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) *Molinio-Arrhenatheretea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R31 Mediterranean tall humid inland grassland](#)

Constant species of EUNIS habitats: [R31 Mediterranean tall humid inland grassland](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Grassland (non-alpine, non-saline)**