Carex maritima

Habitus and growth type

Plant height [m]: **0.08**
Life span: **Perennial**
Life form: **Hemicryptophyte, Geophyte**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **1.13**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.6**
Temperature indicator value: **1.4**
Moisture indicator value: **8.2**
Reaction indicator value: **7**
Nutrient indicator value: **1.8**
Salinity indicator value: **0.6**

Disturbance indicator values

Disturbance frequency: **1.65**
Disturbance frequency (herb layer): **1.88**
Disturbance severity: **0.26**
Disturbance severity (herb layer): **0.26**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.13**
Grazing pressure: **0.19**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **PA (SCH) *Scheuchzerio palustris-Caricetea fuscae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **Q45 Arctic-alpine rich fen**
Constant species of EUNIS habitats: **Q45 Arctic-alpine rich fen**

Broad habitat

Occurrence in broad habitats: **Wetland, Mire, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **7**
Continentality amplitude: **7**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz