

Carex pyrenaica

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **17.97**

Fruit, seed and dispersal

Seed mass [mg]: **0.37**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **1**

Moisture indicator value: **6**

Reaction indicator value: **2.5**

Nutrient indicator value: **2.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.86**

Disturbance frequency (herb layer): **1.86**

Disturbance severity: **0.27**

Disturbance severity (herb layer): **0.27**

Mowing frequency: **0.03**

Grazing pressure: **0.2**

Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [LA \(HER\) *Salicetea herbaceae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R41 Snow-bed vegetation](#)

Broad habitat

Occurrence in broad habitats: **Alpine-subalpine and arctic grassland**