

Carex remota

Habitus and growth type

Plant height [m]: **0.42**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **25.75**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.41**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Reaction indicator value: **5.9**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.17**

Disturbance frequency (herb layer): **1.41**

Disturbance severity: **0.66**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.04**

Grazing pressure: **0.19**

Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [HA \(POP\) *Alno glutinosae-Populetea albae*](#), [OA \(MON\) *Montio-Cardaminetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T12 *Alnus glutinosa*-*Alnus incana* forest on riparian and mineral soils](#), [T13 Temperate hardwood riparian forest](#)

Constant species of EUNIS habitats: [T12 *Alnus glutinosa*-*Alnus incana* forest on riparian and mineral soils](#), [T13 Temperate hardwood riparian forest](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Wetland, Scrub, Forest, Synanthropic**

Distribution

Continentality amplitude: **5**