Carex remota

Habitus and growth type

Plant height [m]: **0.42**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **25.75**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.41**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Reaction indicator value: **5.9**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.17**
Disturbance frequency (herb layer): **1.41**
Disturbance severity: **0.66**
Disturbance severity (herb layer): **0.28**
Mowing frequency: **0.04**
Grazing pressure: **0.19**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **HA (POP) *Alno glutinosae-Populetea albae*, OA (MON) *Montio-Cardaminetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils, T13 Temperate hardwood riparian forest**

Constant species of EUNIS habitats: **T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils, T13 Temperate hardwood riparian forest**

Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Wetland, Scrub, Forest, Synanthropic**

Distribution

Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz