

# *Carex rostrata*

## Habitus and growth type

Plant height [m]: **0.53**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **15.8**

## Flower

Flowering period: **April-July**

## Fruit, seed and dispersal

Seed mass [mg]: **0.95**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **4.3**

Moisture indicator value: **9.9**

Reaction indicator value: **3.6**

Nutrient indicator value: **2.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.07**

Disturbance frequency (herb layer): **1.43**

Disturbance severity: **0.27**

Disturbance severity (herb layer): **0.21**

Mowing frequency: **0.28**

Grazing pressure: **0.12**

Soil disturbance: **0.14**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [IA \(ALN\) \*Alnetea glutinosae\*](#), [OD \(PHR\) \*Phragmito-Magnocaricetea\*](#), [PA \(SCH\) \*Scheuchzerio palustris-Caricetea fuscae\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [Q22 Poor fen](#), [Q24 Intermediate fen and soft-water spring mire](#), [Q25 Non-calcareous quaking mire](#), [Q42 Extremely rich moss-sedge fen](#), [Q44 Calcareous quaking mire](#)

Constant species of EUNIS habitats: [Q22 Poor fen](#), [Q24 Intermediate fen and soft-water spring mire](#), [Q25 Non-calcareous quaking mire](#), [Q41 Alkaline, calcareous, carbonate-rich small-sedge spring fen](#), [Q42 Extremely rich moss-sedge fen](#), [Q43 Tall-sedge base-rich fen](#), [Q44 Calcareous quaking mire](#), [Q53 Tall-sedge bed](#), [T15 Broadleaved swamp forest on non-acid peat](#), [T16 Broadleaved mire forest on acid peat](#)

### Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Wetland, Mire, Grassland (non-alpine, non-saline), Scrub, Forest**

## Distribution

Continentality amplitude: **8**