

Carex sylvatica

Habitus and growth type

Plant height [m]: **0.51**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **30.9**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **1.75**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Moisture indicator value: **5.7**

Reaction indicator value: **6.6**

Nutrient indicator value: **5.1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.06**

Disturbance frequency (herb layer): **1.34**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.17**

Mowing frequency: **0.03**

Grazing pressure: **0.19**

Soil disturbance: **0.11**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [T17 Fagus forest on non-acid soils](#), [T1E Carpinus and Quercus mesic deciduous forest](#)

Constant species of EUNIS habitats: [S37 Corylus avellana scrub](#), [S38 Temperate forest clearing scrub](#), [T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils](#), [T13 Temperate hardwood riparian forest](#), [T17 Fagus forest on non-acid soils](#), [T1E Carpinus and Quercus mesic deciduous forest](#), [T1F Ravine forest](#), [T32 Temperate mountain Abies forest](#)

Broad habitat

Occurrence in broad habitats: **Spring, Scrub, Forest, Synanthropic**

Distribution

Continentality amplitude: **5**