

# ***Centaurea cineraria***

## **Habitus and growth type**

Plant height [m]: **0.62**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## **Flower**

Flowering period: **May-August**

## **Fruit, seed and dispersal**

Seed mass [mg]: **2.54**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **9**

Moisture indicator value: **3**

Reaction indicator value: **7**

Nutrient indicator value: **6**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.88**

Disturbance frequency (herb layer): **0.91**

Disturbance severity: **0.48**

Disturbance severity (herb layer): **0.39**

Mowing frequency: **0.23**

Grazing pressure: **0.31**

Soil disturbance: **0.1**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [\*\*JB \(CRI\) Crithmo-Staticetea\*\*](#)

### Broad habitat

Occurrence in broad habitats: **Coastal cliff, Sparsely vegetated (incl. rock and scree)**