Centaurea linifolia

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**
Spinescence: **not spinescent**

Fruit, seed and dispersal

Dispersal mode: **Myrmecochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.4**
Temperature indicator value: **8.2**
Moisture indicator value: **2.6**
Reaction indicator value: **7.1**
Nutrient indicator value: **2.9**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.4**
Disturbance frequency (herb layer): **1.56**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.21**
Mowing frequency: **0.02**
Grazing pressure: **0.37**
Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FB (ROS) *Ononido-Rosmarinetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S61 Western basiphilous garrigue**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz