

# ***Centaurea spinosa***

## **Habitus and growth type**

Life span: **Perennial**

Life form: **Chamaephyte**

## **Fruit, seed and dispersal**

Seed mass [mg]: **2.23**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Dry**

Salinity relationship: **Slightly saline or brackish**

### **Ellenberg-type indicator values**

Light indicator value: **8**

Temperature indicator value: **11**

Moisture indicator value: **4**

Reaction indicator value: **8**

Nutrient indicator value: **5**

Salinity indicator value: **3**

### **Disturbance indicator values**

Disturbance frequency: **1.22**

Disturbance frequency (herb layer): **1.67**

Disturbance severity: **0.36**

Disturbance severity (herb layer): **0.31**

Mowing frequency: **0.01**

Grazing pressure: **0.19**

Soil disturbance: **0.41**

## **Habitat and sociology**

### **Syntaxon**

Diagnostic species of phytosociological classes: [FB \(ROS\) Ononido-Rosmarinetea](#), [JB \(CRI\) Crithmo-Staticetea](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Coastal cliff, Scrub**