Centranthus angustifolius

Habitus and growth type

Plant height [m]: **0.5**
Life span: **Perennial**
Life form: **Chamaephyte, Hemicryptophyte**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **2.45**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Nutrient relationship: **Dystrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**
Temperature indicator value: **8**
Moisture indicator value: **2**
Nutrient indicator value: **1**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.87**
Disturbance frequency (herb layer): **1.4**
Disturbance severity: **0.52**
Disturbance severity (herb layer): **0.43**
Mowing frequency: **0.14**
Grazing pressure: **0.2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Soil disturbance: **0.28**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **KE (THL) *Thlaspietea rotundifolii***

EUNIS habitat

Diagnostic species of EUNIS habitats: **U72 Unvegetated or sparsely vegetated gravel bar in the Mediterranean region**

Constant species of EUNIS habitats: **U72 Unvegetated or sparsely vegetated gravel bar in the Mediterranean region**

Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz