

Centranthus ruber

Habitus and growth type

Plant height [m]: **0.56**

Life span: **Perennial**

Life form: **Chamaephyte, Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **18.45**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **1.89**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.1**

Moisture indicator value: **3.6**

Reaction indicator value: **8**

Nutrient indicator value: **5**

Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **0.25**

Disturbance frequency (herb layer): **1.33**

Disturbance severity: **0.68**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.08**

Grazing pressure: **0.24**

Soil disturbance: **0.16**

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: **5**

Continentality amplitude: **3**