Cerastium banaticum

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.44**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **7.8**
Temperature indicator value: **5.6**
Moisture indicator value: **4.4**

Disturbance indicator values

Disturbance frequency: **1.43**
Disturbance frequency (herb layer): **1.66**
Disturbance severity: **0.4**
Disturbance severity (herb layer): **0.38**
Mowing frequency: **0.27**
Grazing pressure: **0.3**
Soil disturbance: **0.14**

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz