

Cerastium brachypetalum

Habitus and growth type

Plant height [m]: **0.17**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **21.76**

Flower

Flowering period: **February-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.06**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.8**

Moisture indicator value: **3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.92**

Disturbance frequency (herb layer): **1.84**

Disturbance severity: **0.56**

Disturbance severity (herb layer): **0.48**

Mowing frequency: **0.37**

Grazing pressure: **0.29**

Soil disturbance: **0.3**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [S34 Balkan-Anatolian submontane genistoid scrub](#), [T38 Mediterranean montane Cedrus forest](#)

Constant species of EUNIS habitats: [S34 Balkan-Anatolian submontane genistoid scrub](#), [T38 Mediterranean montane Cedrus forest](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Oromediterranean scrub, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **5**

Continentality amplitude: **4**