Cerastium carinthiacum

Habitus and growth type

Plant height [m]: **0.1**
Life span: **Perennial**
Life form: **Chamaephyte**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.2**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **2.3**
Moisture indicator value: **4.3**
Reaction indicator value: **7.3**
Nutrient indicator value: **3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.05**
Disturbance frequency (herb layer): **1.26**
Disturbance severity: **0.35**
Disturbance severity (herb layer): **0.32**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.01**
Grazing pressure: **0.15**
Soil disturbance: **0.17**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **U26 Temperate high-mountain base-rich scree and moraine**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz