Cirsium brachycephalum

Habitus and growth type

Plant height [m]: **0.85**
Life span: **Biennial or short-lived**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.85**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **7**
Moisture indicator value: **8**
Reaction indicator value: **7.5**
Nutrient indicator value: **5**
Salinity indicator value: **4.7**

Disturbance indicator values

Disturbance frequency: **1.84**
Disturbance frequency (herb layer): **1.84**
Disturbance severity: **0.38**
Disturbance severity (herb layer): **0.38**
Mowing frequency: **1.07**
Grazing pressure: **0.32**
Soil disturbance: **0.13**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DB (FEP) *Festuco-Puccinellietea***

Broad habitat

Occurrence in broad habitats: **Wetland, Saline vegetation**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz