

Cirsium canum

Habitus and growth type

Plant height [m]: **0.8**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **May-September**

Fruit, seed and dispersal

Seed mass [mg]: **2.09**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.9**

Temperature indicator value: **6.3**

Moisture indicator value: **7.6**

Reaction indicator value: **6.7**

Nutrient indicator value: **5.8**

Salinity indicator value: **0.4**

Disturbance indicator values

Disturbance frequency: **1.73**

Disturbance frequency (herb layer): **1.98**

Disturbance severity: **0.52**

Disturbance severity (herb layer): **0.51**

Mowing frequency: **1.28**

Grazing pressure: **0.19**

Soil disturbance: **0.2**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) *Molinio-Arrhenatheretea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R34 Submediterranean moist meadow](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Wetland, Mire, Grassland (non-alpine, non-saline), Saline vegetation**

Distribution

Continentality: **7**

Continentality amplitude: **3**