

# *Cirsium erisithales*

## Habitus and growth type

Plant height [m]: **1.07**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **May-August**

## Fruit, seed and dispersal

Seed mass [mg]: **3.04**

Dispersal mode: **Anemochory**

Dispersal distance class: **3**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **5.8**

Temperature indicator value: **4.3**

Moisture indicator value: **5.8**

Reaction indicator value: **7.8**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.12**

Disturbance frequency (herb layer): **1.15**

Disturbance severity: **0.63**

Disturbance severity (herb layer): **0.17**

Mowing frequency: **0.05**

Grazing pressure: **0.2**

Soil disturbance: **0.11**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CP \(MUG\) \*Roso pendulinae-Pinetea mugo\*](#), [CS \(MUL\) \*Mulgedio-Aconitetea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [S24 Subalpine genistoid scrub of the Amphi-Adriatic region](#)

Constant species of EUNIS habitats: [S24 Subalpine genistoid scrub of the Amphi-Adriatic region](#)

### Broad habitat

Occurrence in broad habitats: **Spring, Alpine-subalpine and arctic grassland, Scrub, Forest**

## Distribution

Continentality: **6**

Continentality amplitude: **3**