

# *Cirsium furiens*

## Habitus and growth type

Life span: **Biennial or short-lived**

Life form: **Hemicryptophyte, Geophyte**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **7**

Moisture indicator value: **4**

Reaction indicator value: **7**

Nutrient indicator value: **7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.9**

Disturbance frequency (herb layer): **1.91**

Disturbance severity: **0.5**

Disturbance severity (herb layer): **0.47**

Mowing frequency: **0.96**

Grazing pressure: **0.29**

Soil disturbance: **0.18**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#), [QF \(ART\) Artemisietea vulgaris](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Synanthropic**