

Cirsium morisianum

Habitus and growth type

Plant height [m]: **0.82**

Life span: **Biennial or short-lived**

Life form: **Hemicryptophyte**

Flower

Flowering period: **May-September**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.5**

Moisture indicator value: **4**

Reaction indicator value: **7**

Nutrient indicator value: **7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.13**

Disturbance frequency (herb layer): **1.78**

Disturbance severity: **0.54**

Disturbance severity (herb layer): **0.5**

Mowing frequency: **0.56**

Grazing pressure: **0.29**

Soil disturbance: **0.18**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QF \(ART\) Artemisietea vulgaris](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**