

Cirsium rivulare

Habitus and growth type

Plant height [m]: **0.55**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **2.4**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.4**

Temperature indicator value: **4.4**

Moisture indicator value: **7.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.36**

Disturbance frequency (herb layer): **1.79**

Disturbance severity: **0.52**

Disturbance severity (herb layer): **0.5**

Mowing frequency: **1**

Grazing pressure: **0.18**

Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) Molinio-Arrhenatheretea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R35 Moist or wet mesotrophic to eutrophic hay meadow](#)

Constant species of EUNIS habitats: [R35 Moist or wet mesotrophic to eutrophic hay meadow](#)

Broad habitat

Occurrence in broad habitats: **Mire, Grassland (non-alpine, non-saline)**

Distribution

Continentality: **5**

Continentality amplitude: **3**