

Cirsium tuberosum

Habitus and growth type

Plant height [m]: **0.54**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

Leaf

Specific leaf area [mm²/mg]: **11.7**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **2.99**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.5**

Temperature indicator value: **5.8**

Reaction indicator value: **7.8**

Nutrient indicator value: **2.7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.71**

Disturbance frequency (herb layer): **1.74**

Disturbance severity: **0.52**

Disturbance severity (herb layer): **0.39**

Mowing frequency: **0.59**

Grazing pressure: **0.26**

Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) Molinio-Arrhenatheretea](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Wetland, Grassland (non-alpine, non-saline)**

Distribution

Continentality: **4**

Continentality amplitude: **2**