Cistus atriplicifolius

Habitus and growth type

Life span: **Perennial**
Life form: **Phanerophyte, Shrub**
Spinescence: **not spinescent**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Disturbance indicator values

Disturbance frequency: **0.29**
Disturbance frequency (herb layer): **1.56**
Disturbance severity: **0.7**
Disturbance severity (herb layer): **0.16**
Mowing frequency: **0**
Grazing pressure: **0.34**
Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FC (LAV) *Cisto-Lavanduletea stoechadis***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S62 Western acidophilous garrigue**

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz