Cistus crispus

Habitus and growth type

Plant height [m]: **0.46**
Life span: **Perennial**
Life form: **Phanerophyte, Shrub**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **9.48**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **0.48**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **8.5**
Moisture indicator value: **3**
Reaction indicator value: **3**
Nutrient indicator value: **3**
Salinity indicator value: **0**

Disturbance indicator values

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency: **0.2**
Disturbance frequency (herb layer): **1.46**
Disturbance severity: **0.66**
Disturbance severity (herb layer): **0.14**
Mowing frequency: **0**
Grazing pressure: **0.3**
Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FC (LAV) *Cisto-Lavanduletea stoechadis***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S62 Western acidophilous garrigue**

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz