

Cistus ocymoides

Habitus and growth type

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.7**

Temperature indicator value: **7.2**

Moisture indicator value: **3.3**

Reaction indicator value: **2.5**

Nutrient indicator value: **1.7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.55**

Disturbance frequency (herb layer): **1.49**

Disturbance severity: **0.51**

Disturbance severity (herb layer): **0.21**

Mowing frequency: **0**

Grazing pressure: **0.34**

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CH \(ULI\) *Calluno-Ulicetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S62 *Western acidophilous garrigue*](#)

Broad habitat

Occurrence in broad habitats: **Heathland, Scrub**