

Cistus psilosepalus

Habitus and growth type

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Fruit, seed and dispersal

Seed mass [mg]:

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **7**

Moisture indicator value: **2**

Reaction indicator value: **2**

Nutrient indicator value: **1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.42**

Disturbance frequency (herb layer): **1.36**

Disturbance severity: **0.48**

Disturbance severity (herb layer): **0.2**

Mowing frequency: **0.01**

Grazing pressure: **0.27**

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CH \(ULI\) Calluno-Ulicetea](#), [FA \(QUI\)](#)

[Quercetea ilicis](#)

Broad habitat

Occurrence in broad habitats: **Heathland, Forest**