Clematis integrifolia

Habitus and growth type

Plant height [m]: **0.43**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **6.9**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.5**
Moisture indicator value: **5.9**
Reaction indicator value: **7**
Nutrient indicator value: **5**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.21**
Disturbance frequency (herb layer): **1.86**
Disturbance severity: **0.51**
Disturbance severity (herb layer): **0.48**
Mowing frequency: **0.77**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.26**
Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CM (MOL) *Molinio-Arrhenatheretea***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz