

Clinopodium graveolens

Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.39**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Temperature indicator value: **11**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.86**

Disturbance frequency (herb layer): **1.67**

Disturbance severity: **0.44**

Disturbance severity (herb layer): **0.37**

Mowing frequency: **0.21**

Grazing pressure: **0.32**

Soil disturbance: **0.19**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [S34 Balkan-Anatolian submontane genistoid scrub](#), [T38 Mediterranean montane Cedrus forest](#), [T3D Mediterranean Cupressaceae forest](#)

Constant species of EUNIS habitats: [S34 Balkan-Anatolian submontane genistoid scrub](#), [T38 Mediterranean montane Cedrus forest](#), [T3D Mediterranean](#)

Cupressaceae forest

Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Synanthropic**