Cochlearia pyrenaica

Habitus and growth type

Plant height [m]: **0.18**
Life span: **Biennial or short-lived, Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **23.38**

Fruit, seed and dispersal

Seed mass [mg]: **1.09**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.7**
Temperature indicator value: **3.7**
Moisture indicator value: **8**
Reaction indicator value: **7.7**
Nutrient indicator value: **3.3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.3**
Disturbance frequency (herb layer): **1.31**
Disturbance severity: **0.35**
Disturbance severity (herb layer): **0.35**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.29**
Grazing pressure: **0.13**
Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **OA (MON) *Montio-Cardaminetea***

Broad habitat

Occurrence in broad habitats: **Spring**

Distribution

Continentality: **4**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz