

# *Coincya monensis*

## Habitus and growth type

Plant height [m]: **0.44**

Life span: **Biennial or short-lived**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **16.1**

## Flower

Flowering period: **April-August**

## Fruit, seed and dispersal

Seed mass [mg]: **1.07**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **9**

Moisture indicator value: **3.8**

Nutrient indicator value: **2.8**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.34**

Disturbance frequency (herb layer): **1.52**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.45**

Mowing frequency: **0.27**

Grazing pressure: **0.22**

Soil disturbance: **0.26**

### **Habitat and sociology**

#### Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree), Synanthropic**

### **Distribution**

Continentality: **3**

Continentality amplitude: **3**