

# *Convolvulus cantabrica*

## Habitus and growth type

Plant height [m]: **0.35**

Life span: **Perennial**

Life form: **Chamaephyte, Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **19.49**

## Flower

Flowering period: **April-October**

## Fruit, seed and dispersal

Seed mass [mg]: **5.4**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.7**

Temperature indicator value: **7.6**

Moisture indicator value: **2.4**

Reaction indicator value: **7.5**

Nutrient indicator value: **2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.87**

Disturbance frequency (herb layer): **1.78**

Disturbance severity: **0.59**

Disturbance severity (herb layer): **0.48**

Mowing frequency: **0.36**

Grazing pressure: **0.34**

Soil disturbance: **0.23**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#), [FF \(LYG\) Lygeo sparti-Stipetea tenacissimae](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#)

Constant species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#),  
[R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**