

Coriaria myrtifolia

Habitus and growth type

Plant height [m]:

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Flower

Flowering period: **February-April**

Fruit, seed and dispersal

Seed mass [mg]: **9.9**

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with frankia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **7.3**

Moisture indicator value: **5.5**

Reaction indicator value: **6**

Nutrient indicator value: **3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.09**

Disturbance frequency (herb layer): **1.35**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.14**

Mowing frequency: **0.01**

Grazing pressure: **0.23**

Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CD \(RHA\) Crataego-Prunetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S53 Spartium junceum scrub](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Sparsely vegetated (incl. rock and scree)**