

# *Corispermum intermedium*

## Habitus and growth type

Plant height [m]: **0.32**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **21.24**

## Flower

Flowering period: **June-September**

## Fruit, seed and dispersal

Seed mass [mg]: **1.99**

Dispersal mode: **Anemochory**

Dispersal distance class: **4**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **7**

Moisture indicator value: **2.5**

Reaction indicator value: **7**

Nutrient indicator value: **6**

Salinity indicator value: **1.5**

### Disturbance indicator values

Disturbance frequency: **1.73**

Disturbance frequency (herb layer): **1.99**  
Disturbance severity: **0.75**  
Disturbance severity (herb layer): **0.75**  
Mowing frequency: **0.94**  
Grazing pressure: **0.14**  
Soil disturbance: **0.6**

### **Habitat and sociology**

#### Syntaxon

Diagnostic species of phytosociological classes: [QD \(DIG\) \*Digitario sanguinalis-Eragrostietea minoris\*](#)

#### Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Wetland, Synanthropic**

### **Distribution**

Continentality: **6**  
Continentality amplitude: **1**