

# *Cotoneaster tomentosus*

## Habitus and growth type

Plant height [m]:

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

## Flower

Flowering period: **March-May**

## Fruit, seed and dispersal

Seed mass [mg]: **89.29**

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**

Moisture indicator value: **3.3**

Reaction indicator value: **8.8**

Nutrient indicator value: **2.3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.11**

Disturbance frequency (herb layer): **1.21**

Disturbance severity: **0.64**

Disturbance severity (herb layer): **0.14**

Mowing frequency: **0.02**

Grazing pressure: **0.21**

Soil disturbance: **0.1**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CD \(RHA\) Crataego-Prunetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T36 Temperate and submediterranean montane Pinus sylvestris-Pinus nigra forest](#)

Constant species of EUNIS habitats: [T36 Temperate and submediterranean montane Pinus sylvestris-Pinus nigra forest](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **5**

Continentality amplitude: **3**