

# *Crepis fraasii*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **3**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **3.7**

Temperature indicator value: **10**

Moisture indicator value: **5.7**

Reaction indicator value: **7.5**

Nutrient indicator value: **4.3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.1**

Disturbance frequency (herb layer): **1.46**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.23**

Mowing frequency: **0.02**

Grazing pressure: **0.29**

Soil disturbance: **0.13**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **[CB \(PUB\) Quercetea pubescentis](#)**

EUNIS habitat

Diagnostic species of EUNIS habitats: [T3D Mediterranean Cupressaceae forest](#)

Constant species of EUNIS habitats: [T3D Mediterranean Cupressaceae forest](#)

Broad habitat

Occurrence in broad habitats: **Forest**