

Crepis pusilla

Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **11**

Moisture indicator value: **5**

Reaction indicator value: **8**

Nutrient indicator value: **6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.4**

Disturbance frequency (herb layer): **1.94**

Disturbance severity: **0.82**

Disturbance severity (herb layer): **0.8**

Mowing frequency: **0.25**

Grazing pressure: **0.5**

Soil disturbance: **0.47**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [OC \(ISO\) Isoëto-Nanojuncetea](#), [QE](#)

(POL) *Polygono-Poetea annuae*

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1D Mediterranean closely grazed dry grassland**

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Synanthropic**