Crepis setosa

Habitus and growth type

Plant height [m]: **0.32**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **April-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.18**
Dispersal mode: **Anemochory, Anthropochory**
Dispersal distance class: **5, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **7.5**
Moisture indicator value: **3.8**
Reaction indicator value: **7.4**
Nutrient indicator value: **4.4**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.57**
Disturbance frequency (herb layer): **2.01**
Disturbance severity: **0.66**
Disturbance severity (herb layer): **0.64**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.72**
Grazing pressure: **0.26**
Soil disturbance: **0.34**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QB (SIS) *Sisymbrietea***, **QF (ART) *Artemisietea vulgaris***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R34 Submediterranean moist meadow**, **S34 Balkan-Anatolian submontane genistoid scrub**

Constant species of EUNIS habitats: **R34 Submediterranean moist meadow**, **S34 Balkan-Anatolian submontane genistoid scrub**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**

Distribution

Continentality: **5**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz