

# *Crocus veluchensis*

## Habitus and growth type

Life span: **Perennial**

Life form: **Geophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Ellenberg-type indicator values

Moisture indicator value: **7**

Disturbance indicator values

Disturbance frequency: **0.82**

Disturbance frequency (herb layer): **1.39**

Disturbance severity: **0.36**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.1**

Grazing pressure: **0.24**

Soil disturbance: **0.09**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CT \(TRI\) \*Juncetea trifidi\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [Q23 Relict mire of Mediterranean mountains](#), [R1K Balkan and Anatolian oromediterranean dry grassland](#), [T39 Mediterranean and Balkan subalpine \*Pinus heldreichii\*-\*Pinus peuce\* forest](#)

Constant species of EUNIS habitats: [T39 Mediterranean and Balkan subalpine \*Pinus heldreichii\*-\*Pinus peuce\* forest](#)

## Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Forest**