

Crypsis alopecuroides

Habitus and growth type

Plant height [m]: **0.15**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **June–October**

Fruit, seed and dispersal

Seed mass [mg]: **0.2**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Ellenberg-type indicator values

Light indicator value: **8.3**

Moisture indicator value: **6.8**

Reaction indicator value: **7.5**

Salinity indicator value: **3.8**

Disturbance indicator values

Disturbance frequency: **1.7**

Disturbance frequency (herb layer): **1.85**

Disturbance severity: **0.65**

Disturbance severity (herb layer): **0.65**

Mowing frequency: **0.25**

Grazing pressure: **0.24**

Soil disturbance: **0.33**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**OC \(ISO\) Isoëto-Nanojuncetea**](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**Q63 Periodically exposed saline shore with pioneer or ephemeral vegetation**](#)

Dominant species of EUNIS habitats: [**Q63 Periodically exposed saline shore with pioneer or ephemeral vegetation**](#)

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Wetland, Saline vegetation, Sparsely vegetated (incl. rock and scree)**