

Cuscuta campestris

Habitus and growth type

Plant height [m]: **0.3**

Life span: **Annual**

Life form: **Therophyte, Herbaceous liana**

Flower

Flowering period: **April-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.71**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **parasitic vine**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Neophyte**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.2**

Moisture indicator value: **5.1**

Reaction indicator value: **7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.38**

Disturbance frequency (herb layer): **1.81**

Disturbance severity: **0.71**

Disturbance severity (herb layer): **0.69**

Mowing frequency: **0.41**

Grazing pressure: **0.18**

Soil disturbance: **0.5**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QA \(PAR\) *Papaveretea rhoeadis*](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Grassland (non-alpine, non-saline), Synanthropic**

Distribution

Continentality: **4**

Continentality amplitude: **4**