

# *Cyclamen hederifolium*

## Habitus and growth type

Plant height [m]: **0.1**

Life span: **Perennial**

Life form: **Geophyte**

## Flower

Flowering period: **July-October**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **3.3**

Temperature indicator value: **8.5**

Moisture indicator value: **4.7**

Reaction indicator value: **6.3**

Nutrient indicator value: **4.7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.03**

Disturbance frequency (herb layer): **1.31**

Disturbance severity: **0.7**

Disturbance severity (herb layer): **0.12**

Mowing frequency: **0**

Grazing pressure: **0.22**

Soil disturbance: **0.1**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CB \(PUB\) \*Quercetea pubescentis\*](#), [FA \(QUI\) \*Quercetea ilicis\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T1G \*Alnus cordata\* forest](#), [T22 Mainland laurophyllous forest](#)

Constant species of EUNIS habitats: [T1G \*Alnus cordata\* forest](#), [T22 Mainland laurophyllous forest](#)

Broad habitat

Occurrence in broad habitats: **Forest**