Cytisus blockianus

Habitus and growth type

Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Fruit, seed and dispersal

Dispersal mode: **Myrmecochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **7.5**
Temperature indicator value: **4.8**
Moisture indicator value: **4.8**

Disturbance indicator values

Disturbance frequency: **1.68**
Disturbance frequency (herb layer): **1.68**
Disturbance severity: **0.47**
Disturbance severity (herb layer): **0.47**
Mowing frequency: **0.71**
Grazing pressure: **0.31**
Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz