Cytisus multiflorus

Habitus and growth type

Plant height [m]: **2.1**
Life span: **Perennial**
Life form: **Phanerophyte, Shrub**
Spinescence: **not spinescent**

Fruit, seed and dispersal

Seed mass [mg]: **5.45**
Dispersal mode: **Myrmecochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **8**
Moisture indicator value: **3.5**
Reaction indicator value: **2.5**
Nutrient indicator value: **2.5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.97**
Disturbance frequency (herb layer): **1.35**
Disturbance severity: **0.29**
Disturbance severity (herb layer): **0.22**
Mowing frequency: **0.02**
Grazing pressure: **0.3**
Soil disturbance: **0.12**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FE (CYT) *Cytisetea scopario-striati***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S33 Lowland to montane temperate and submediterranean genistoid scrub**

Dominant species of EUNIS habitats: **S33 Lowland to montane temperate and submediterranean genistoid scrub**

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz