

# ***Daboecia cantabrica***

## **Habitus and growth type**

Plant height [m]: **0.5**

Life span: **Perennial**

Life form: **Chamaephyte**

## **Fruit, seed and dispersal**

Seed mass [mg]: **0.11**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **7**

Moisture indicator value: **5**

Reaction indicator value: **2.7**

Nutrient indicator value: **1.7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.32**

Disturbance frequency (herb layer): **1.36**

Disturbance severity: **0.58**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.05**

Grazing pressure: **0.26**

Soil disturbance: **0.1**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CH \(ULI\) Calluno-Ulicetea](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T29 Broadleaved evergreen plantation of non site-native trees](#)

Constant species of EUNIS habitats: [T29 Broadleaved evergreen plantation of non site-native trees](#)

### Broad habitat

Occurrence in broad habitats: **Heathland, Forest**