

# ***Dianthus armeria***

## **Habitus and growth type**

Plant height [m]: **0.34**

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **15.26**

## **Flower**

Flowering period: **April-August**

## **Fruit, seed and dispersal**

Seed mass [mg]: **0.29**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **7**

Moisture indicator value: **4.2**

Nutrient indicator value: **2.7**

Salinity indicator value: **0**

### **Disturbance indicator values**

Disturbance frequency: **0.69**

Disturbance frequency (herb layer): **1.76**

Disturbance severity: **0.55**

Disturbance severity (herb layer): **0.43**

Mowing frequency: **0.57**

Grazing pressure: **0.26**

Soil disturbance: **0.25**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [CH \(ULI\) \*Calluno-Ulicetea\*](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Heathland, Scrub, Forest**

## **Distribution**

Continentality: **4**

Continentality amplitude: **4**