Dianthus pungens

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**
Spinescence: **not spinescent**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **4**
Moisture indicator value: **1**
Reaction indicator value: **7**
Nutrient indicator value: **1**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.64**
Disturbance frequency (herb layer): **1.53**
Disturbance severity: **0.57**
Disturbance severity (herb layer): **0.36**
Mowing frequency: **0.21**
Grazing pressure: **0.4**
Soil disturbance: **0.17**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1H Iberian oromediterranean basiphilous dry grassland**, **S73 Western Mediterranean mountain hedgehog-heath**
Constant species of EUNIS habitats: **R1H Iberian oromediterranean basiphilous dry**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

grassland, S73 Western Mediterranean mountain hedgehog-heath

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Oromediterranean scrub, Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz