

# ***Dianthus serotinus***

## **Habitus and growth type**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## **Fruit, seed and dispersal**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **6.4**

Moisture indicator value: **3.5**

Reaction indicator value: **8**

Nutrient indicator value: **1**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.63**

Disturbance frequency (herb layer): **1.57**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.29**

Mowing frequency: **0.27**

Grazing pressure: **0.31**

Soil disturbance: **0.27**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CJ \(COR\) Koelerio-Coryneporetea  
canescentis](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**