

Dictamnus albus

Habitus and growth type

Plant height [m]: **0.7**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **20.94**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **14.84**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.5**

Temperature indicator value: **7.2**

Moisture indicator value: **3.2**

Reaction indicator value: **7.8**

Nutrient indicator value: **2.2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.17**

Disturbance frequency (herb layer): **1.39**

Disturbance severity: **0.61**

Disturbance severity (herb layer): **0.2**

Mowing frequency: **0.1**

Grazing pressure: **0.23**

Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CB \(PUB\) Quercetea pubescentis](#), [CL \(GER\) Trifolio-Geranietea sanguinei](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R51 Thermophilous forest fringe of base-rich soils](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **7**

Continentality amplitude: **4**