

# *Diplotaxis viminea*

## Habitus and growth type

Plant height [m]: **0.12**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **January-November**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.8**

Moisture indicator value: **3.3**

Reaction indicator value: **7.3**

Nutrient indicator value: **6**

Salinity indicator value: **0.3**

### Disturbance indicator values

Disturbance frequency: **2.04**

Disturbance frequency (herb layer): **2.04**

Disturbance severity: **0.82**

Disturbance severity (herb layer): **0.82**

Mowing frequency: **0.96**

Grazing pressure: **0.22**

Soil disturbance: **0.58**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) Chenopodietea](#)

### Broad habitat

Occurrence in broad habitats: **Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

## Distribution

Continentality amplitude: 5